

MATHEMATICAL INSIGHTS INTO FRACTIONAL POWER SERIES SOLUTIONS

Haojun Li

School of Mathematics and Statistics, Zhaoqing University, Guangdong, China

Abstract: In this paper, we obtain the exact solutions of two fractional power series. A new multiplication of fractional power series and Jumarie type of Riemann-Liouville (R-L) fractional calculus play important roles in this article. In fact, our results are generalizations of ordinary calculus results.

Keyword: Exact solutions, fractional power series, new multiplication, Jumarie type of R-L fractional calculus.

I. INTRODUCTION

During the 18th and 19th centuries, there were many famous scientists such as Euler, Laplace, Fourier, Abel, Liouville, Grunwald, Letnikov, Riemann, and some others who reported interesting results within fractional calculus. In recent years, fractional calculus has become an increasingly popular research area due to its effective applications in different scientific fields such as economics, engineering, dynamics, biology, control theory, mechanics, chaos theory, and so on [1-10].

However, the definition of fractional derivative is not unique. Common definitions include Riemann-Liouville (R-L) fractional derivative, Caputo fractional derivative, Grunwald-Letnikov (G-L) fractional derivative, and Jumarie's modified R-L fractional derivative [11-14]. Because Jumarie's modified R-L fractional derivative helps avoid non-zero fractional derivative of constant functions, it is easier to use this definition to associate fractional calculus with classical calculus.

In this paper, we find the exact solutions of the following two α -fractional power series:

$$\sum_{n=1}^{\infty} \frac{(4n)!}{\Gamma((4n+1)\alpha+1)} x^{(4n+1)\alpha}, \quad (1)$$

and

$$\sum_{n=0}^{\infty} \frac{1}{\Gamma(4n\alpha+1)} x^{4n\alpha}. \quad (2)$$

where $0 < \alpha \leq 1$. Jumarie type of R-L fractional calculus and a new multiplication of fractional power series play important roles in this paper. And our results are generalizations of the results in classical calculus.

II. PRELIMINARIES

At first, we introduce the fractional derivative used in this paper and its properties.

Definition 2.1 ([15]): Let $0 < \alpha \leq 1$, and x_0 be a real number. The Jumarie type of Riemann-Liouville (R-L) α -fractional derivative is defined by

$$({}_0 D^\alpha)[f(x)] = \frac{1}{x} \frac{d}{dx} \int_{x_0}^x \frac{f(t) - f(x_0)}{(x-t)^{\alpha}} dt. \quad (3)$$

And the Jumarie type of Riemann-Liouville α -fractional integral is defined by

$({}_0 I^\alpha)[f(x)] = \frac{1}{\Gamma(\alpha)} \int_{x_0}^x (x-t)^{\alpha-1} f(t) dt$, (4) where $\Gamma(\cdot)$ is the gamma function. On the other hand, for any positive integer p , we define $({}_0 D_x^\alpha)^p [f(x)] = ({}_0 D_x^\alpha)({}_0 D_x^\alpha) \cdots ({}_0 D_x^\alpha)[f(x)]$, the p -th order α -fractional derivative of $f(x)$.

Proposition 2.2 ([16]): If α, β, x_0, C are real numbers and $\beta \geq \alpha > 0$, then

$$({}_0 D_x^\alpha)[x^\beta] = \Gamma(\beta - \alpha + 1) x^{\beta - \alpha}, \quad (5)$$

and

$$({}_0 D_x^\alpha)[C] = 0. \quad (6)$$

Next, we introduce the definition of fractional power series.

Definition 2.3: Suppose that x and a_n are real numbers for all n , and $0 < \alpha \leq 1$. If the function $f_\alpha: [a, b] \rightarrow R$ can be expressed as $f_\alpha(x^\alpha) = \sum_{n=0}^{\infty} \frac{a_n}{\Gamma(n\alpha+1)} x^{n\alpha}$, then we say that $f_\alpha(x^\alpha)$ is a α -fractional power series.

In the following, a new multiplication of fractional power series is introduced.

Definition 2.4 ([17]): If $0 < \alpha \leq 1$. If $f_\alpha(x^\alpha)$ and $g_\alpha(x^\alpha)$ are two α -fractional power series,

$$f_\alpha(x^\alpha) = \sum_{n=0}^{\infty} \frac{a_n}{\Gamma(n\alpha+1)} x^{n\alpha} = \sum_{n=0}^{\infty} \frac{1}{n!} x^{n\alpha} \quad a_n n! (\Gamma(\alpha+1)) \otimes n, \quad (7)$$

$\otimes n$

$$g_\alpha(x^\alpha) = \sum_{n=0}^{\infty} \frac{b_n}{\Gamma(n\alpha+1)} x^{n\alpha} = \sum_{n=0}^{\infty} \frac{b_n}{n!} \left(\frac{1}{\Gamma(\alpha+1)} x^{n\alpha} \right) \quad (8) \text{ Then we define}$$

$$f_\alpha(x^\alpha) \otimes g_\alpha(x^\alpha) = \sum_{n=0}^{\infty} \frac{1}{\Gamma(n\alpha+1)} x^{n\alpha} \otimes \sum_{n=0}^{\infty} \frac{b_n}{\Gamma(n\alpha+1)} a_n b x^{n\alpha}$$

$$\Gamma(n\alpha+1)$$

$$= \sum_{n=0}^{\infty} \frac{1}{\Gamma(n\alpha+1)} \left(\sum_{m=0}^n \frac{a_m b_{n-m}}{m!} \right) x^{n\alpha}. \quad (9)$$

$$a_{n-m} b_m) x$$

Equivalently,

$$f_\alpha(x^\alpha) \otimes g_\alpha(x^\alpha)$$

$\otimes n \otimes n$

$$= \sum_{n=0}^{\infty} \frac{a_n}{n!} \left(\frac{1}{\Gamma(\alpha+1)} x^{n\alpha} \right) \otimes \sum_{n=0}^{\infty} \frac{b_n}{n!} \left(\frac{1}{\Gamma(\alpha+1)} x^{n\alpha} \right) = \sum_{n=0}^{\infty} \frac{1}{n!} \left(\sum_{m=0}^n \frac{a_m b_{n-m}}{m!} \right) x^{n\alpha} \quad (10)$$

Definition 2.5 ([18]): Assume that $0 < \alpha \leq 1$, and $f_\alpha(x^\alpha), g_\alpha(x^\alpha)$ are α -fractional power series,

$\otimes n$

$$f_\alpha(x^\alpha) = \sum_{n=0}^{\infty} \frac{a_n}{\Gamma(n\alpha+1)} x^{n\alpha} = \sum_{n=0}^{\infty} \frac{a_n}{n!} \left(\frac{1}{\Gamma(\alpha+1)} x^{n\alpha} \right) \quad (11)$$

$\otimes n$

$$g_\alpha(x^\alpha) = \sum_{n=0}^{\infty} \frac{b_n}{\Gamma(n\alpha+1)} x^{n\alpha} = \sum_{n=0}^{\infty} \frac{b_n}{n!} \left(\frac{1}{\Gamma(\alpha+1)} x^{n\alpha} \right) \quad (12) \text{ The compositions of } f_\alpha(x^\alpha) \text{ and } g_\alpha(x^\alpha) \text{ are defined by}$$

$\otimes n$

$$(f \circ g)(x^\alpha) = f(g(x^\alpha)) = \sum_{n=0}^{\infty} \frac{a_n}{n!} (g(x^\alpha))^n, \quad (13)$$

and

$$(g \circ f)(x^\alpha) = g(f(x^\alpha)) = \sum_{n=0}^{\infty} \frac{b_n}{n!} (f(x^\alpha))^n. \quad (14)$$

Definition 2.6 ([19]): If $0 < \alpha \leq 1$, and x is a real number. The α -fractional exponential function is defined by

$$E_\alpha(x^\alpha) = \sum_{n=0}^{\infty} \frac{x^n}{n!} = \sum_{n=0}^{\infty} \frac{1}{n!} x^n = \left(\frac{1}{\Gamma(\alpha+1)} x^\alpha \right)_{\Gamma(\alpha+1)}. \quad (15)$$

And the α -fractional logarithmic function $L_\alpha(x^\alpha)$ is the inverse function of $E_\alpha(x^\alpha)$.

Definition 2.7 ([20]): The α -fractional cosine and sine function are defined respectively as follows:

$$\cos_\alpha(x^\alpha) = \sum_{n=0}^{\infty} \frac{(-1)^n}{(2n)! \Gamma(\alpha+1)} x^{2n\alpha} = \sum_{n=0}^{\infty} \frac{(-1)^n}{(2n)! \Gamma(\alpha+1)} \left(\frac{1}{\Gamma(\alpha+1)} x^\alpha \right)^{2n}. \quad (16)$$

and

$$\sin_\alpha(x^\alpha) = \sum_{n=0}^{\infty} \frac{(-1)^n}{(2n+1)! \Gamma(\alpha+1)} x^{(2n+1)\alpha} = \sum_{n=0}^{\infty} \frac{(-1)^n}{(2n+1)! \Gamma(\alpha+1)} \left(\frac{1}{\Gamma(\alpha+1)} x^\alpha \right)^{2n+1}. \quad (17)$$

III. EXAMPLES

In this section, we find the exact solutions of two fractional power series.

Example 3.1: Suppose that $0 < \alpha \leq 1$. Find the α -fractional power series

$$\sum_{n=1}^{\infty} \frac{(4n)!}{\Gamma((4n+1)\alpha+1)} x^{(4n+1)\alpha}, \quad (18)$$

where $\frac{1}{\Gamma(\alpha+1)} - 1 < x^\alpha < 1$, and $\left(\frac{1}{\Gamma(\alpha+1)} x^\alpha \right)^4 < 1$.

$\Gamma(\alpha+1)$

$$\text{Solution } \sum_{n=1}^{\infty} \frac{(4n)!}{\Gamma((4n+1)\alpha+1)} x^{(4n+1)\alpha}$$

$$\otimes (4n+1)$$

$$= \sum_{n=1}^{\infty} \frac{1}{4n+1} \left(\frac{1}{\Gamma(\alpha+1)} x^\alpha \right)$$

$$\otimes 4n$$

$$= \sum_{n=1}^{\infty} \left({}_0 I_x^\alpha \right) \left[\left(\frac{1}{\Gamma(\alpha+1)} x^\alpha \right) \right]$$

$$\otimes 4n$$

$$= \left({}_0 I_x^\alpha \right) \left[\sum_{n=1}^{\infty} \left(\frac{1}{\Gamma(\alpha+1)} x^\alpha \right) \right]$$

$$\otimes 4 \otimes 4n$$

$$= \left({}_0 I_x^\alpha \right) \left[\left(\frac{1}{\Gamma(\alpha+1)} x^\alpha \right) \otimes \sum_{n=0}^{\infty} \left(\frac{1}{\Gamma(\alpha+1)} x^\alpha \right) \right]$$

$$\Gamma(\alpha+1)$$

$$= \left({}_0 I_x^\alpha \right) \left[\left(\frac{1}{\Gamma(\alpha+1)} x^\alpha \right) \otimes \left[1 - \left(\frac{1}{\Gamma(\alpha+1)} x^\alpha \right) \right] \right]$$

$$\otimes 4 \otimes -1$$

$$= \left({}_0 I_x^\alpha \right) \left[-1 + \left[1 - \left(\frac{1}{\Gamma(\alpha+1)} x^\alpha \right) \right] \right]$$

$$\otimes 2 \otimes -1 \otimes 2 \otimes -1$$

$$= \left({}_0 I_x^\alpha \right) \left[-1 + \frac{1}{2} \left[1 + \left(\frac{1}{\Gamma(\alpha+1)} x^\alpha \right) \right] + \frac{1}{2} \left[1 - \left(\frac{1}{\Gamma(\alpha+1)} x^\alpha \right) \right] \right]$$

$$\begin{aligned}
 &= 12\alpha) \left[[1 - \frac{1}{\Gamma(\alpha+1)} - \frac{1}{\Gamma(\alpha+1)} (x^\alpha) \otimes 2] \otimes -1 \right] + 1 (0I x^\alpha) \left[[1 + \right. \\
 &(x^\alpha) \otimes 2] \otimes -1] - (0I x^\alpha) [1] \\
 &(0I x^2 \\
 &- \left(\left(1 + \frac{1}{\Gamma(\alpha+1)} 14\alpha \right) - \frac{1}{\Gamma(\alpha+1)} \alpha \otimes \frac{1}{\Gamma(\alpha+1)} (1 - \frac{1}{\Gamma(\alpha+1)} - x) \otimes -1 \right) + \\
 &\arctan_\alpha(x^\alpha) \\
 &= L n_\alpha x^{\frac{1}{\Gamma(\alpha+1)}}
 \end{aligned}
 \tag{19}$$

Example 3.2: Let $0 < \alpha \leq 1$. Find the α -fractional power series

$$\sum_{n=0}^{\infty} \frac{1}{\Gamma(4n\alpha+1)} x^{4n\alpha}. \tag{20}$$

Solution Let $y_\alpha(x^\alpha) = \sum_{n=0}^{\infty} \frac{1}{\Gamma(4n\alpha+1)} x^{4n\alpha}$, then the α -fractional derivatives of $y_\alpha(x^\alpha)$

$$({}_0D_x^\alpha)[y_\alpha(x^\alpha)] = \sum_{n=1}^{\infty} \frac{1}{\Gamma((4n-1)\alpha+1)} x^{(4n-1)\alpha}, \tag{21}$$

$$({}_0D_x^\alpha)^2[y_\alpha(x^\alpha)] = \sum_{n=1}^{\infty} \frac{1}{\Gamma((4n-2)\alpha+1)} x^{(4n-2)\alpha}, \tag{22}$$

$$({}_0D_x^\alpha)^3[y_\alpha(x^\alpha)] = \sum_{n=1}^{\infty} \frac{1}{\Gamma((4n-3)\alpha+1)} x^{(4n-3)\alpha}, \tag{23} \quad ({}_0D_x^\alpha)^4[y_\alpha(x^\alpha)] = \sum_{n=1}^{\infty} \frac{1}{\Gamma((4n-4)\alpha+1)} x^{(4n-4)\alpha} = \sum_{n=0}^{\infty} \frac{1}{\Gamma(4n\alpha+1)} x^{4n\alpha}. \tag{24}$$

Therefore,

$$({}_0D_x^\alpha)^4[y_\alpha(x^\alpha)] - y_\alpha(x^\alpha) = 0, \tag{25} \text{ and}$$

$$y_\alpha(0) = 1, ({}_0D_x^\alpha)[y_\alpha(x^\alpha)](0) = ({}_0D_x^\alpha)^2[y_\alpha(x^\alpha)](0) = ({}_0D_x^\alpha)^3[y_\alpha(x^\alpha)](0) = 0. \tag{26}$$

This is a 4-th order linear α -fractional differential equation, and we can obtain the general solution is

$$y_\alpha(x^\alpha) = C_1 E_\alpha(x^\alpha) + C_2 E_\alpha(-x^\alpha) + C_3 \cos_\alpha(x^\alpha) + C_4 \sin_\alpha(x^\alpha), \tag{27} \text{ where } C_1, C_2, C_3, C_4 \text{ are constants.}$$

Using initial value conditions yields

$$C_1 = C_2 = 1, C_3 = \frac{1}{2}, \text{ and } C_4 = 0. \tag{28}$$

Thus,

$$y_\alpha(x^\alpha) = {}_1_4 E_\alpha(x^\alpha) + {}_4_1 E_\alpha(-x^\alpha) + \frac{1}{2} \cos_\alpha(x^\alpha). \tag{29}$$

That is,

$$\sum_{n=0}^{\infty} \frac{1}{\Gamma(4n\alpha+1)} x^{4n\alpha} = \frac{1}{4} E_\alpha(x^\alpha) + \frac{1}{4} E_\alpha(-x^\alpha) + \frac{1}{2} \cos_\alpha(x^\alpha). \tag{30}$$

IV. CONCLUSION

In this paper, we find the exact solutions of two fractional power series. Jumarie's modified R-L fractional calculus and a new multiplication of fractional power series play important roles in this article. In fact, our results are generalizations of the results in traditional calculus. In the future, we will expand our research fields to engineering mathematics and fractional differential equations.

REFERENCES

- F. Duarte and J. A. T. Machado, Chaotic phenomena and fractional-order dynamics in the trajectory control of redundant manipulators, Nonlinear Dynamics, vol. 29, no. 1-4, pp. 315-342, 2002.
- H. A. Fallahgoul, S. M. Focardi and F. J. Fabozzi, Fractional calculus and fractional processes with applications to financial economics, theory and application, Elsevier Science and Technology, 2016.
- R. Magin, Fractional calculus in bioengineering, part 1, Critical Reviews in Biomedical Engineering, vol. 32, no.1, pp.1-104, 2004.
- Mohd. Farman Ali, Manoj Sharma, Renu Jain, An application of fractional calculus in electrical engineering, Advanced Engineering Technology and Application, vol. 5, no. 2, pp. 41-45, 2016.

- R. Hilfer (Ed.), Applications of Fractional Calculus in Physics, WSPC, Singapore, 2000.
- Teodor M. Atanacković, Stevan Pilipović, Bogoljub Stanković, Dušan Zorica, Fractional Calculus with Applications in Mechanics: Vibrations and Diffusion Processes, John Wiley & Sons, Inc., 2014.
- B. M. Vinagre and Yang Quan Chen, Fractional calculus applications in automatic control and robotics, 41st IEEE Conference on decision and control Tutorial Workshop #2, Las Vegas, December 2002.
- R. L. Bagley and P. J. Torvik, A theoretical basis for the application of fractional calculus to viscoelasticity, Journal of Rheology, vol. 27, no. 3, pp. 201-210, 1983.
- J. Sabatier, OP Agrawal, JA Tenreiro machado, Advances in fractional calculus, Theoretical developments and applications in physics and engineering, vol. 736 Springer; 2007.
- C. -H. Yu, A new insight into fractional logistic equation, International Journal of Engineering Research and Reviews, vol. 9, no. 2, pp.13-17, 2021.
- I. Podlubny, Fractional Differential Equations, Academic Press, San Diego, Calif, USA, 1999.
- S. Das, Functional Fractional Calculus, 2nd Edition, Springer-Verlag, 2011.
- K. B. Oldham, J. Spanier, The Fractional Calculus; Academic Press: New York, NY, USA, 1974.
- K. S. Miller, B. Ross, An Introduction to the Fractional Calculus and Fractional Differential Equations; John Wiley and Sons, Inc.: New York, NY, USA, 1993.
- C. -H. Yu, Application of differentiation under fractional integral sign, International Journal of Mathematics and Physical Sciences Research, vol. 10, no. 2, pp. 40-46, 2022.
- U. Ghosh, S. Sengupta, S. Sarkar and S. Das, Analytic solution of linear fractional differential equation with Jumarie derivative in term of Mittag-Leffler function, American Journal of Mathematical Analysis, vol. 3, no. 2, pp. 32-38, 2015.
- C. -H. Yu, Fractional integral curves of some fractional differential equations, International Journal of Electrical and Electronics Research, vol. 10, no. 4, pp. 17-22, 2022.
- C. -H. Yu, Research on fractional exponential function and logarithmic function, International Journal of Novel Research in Interdisciplinary Studies, vol. 9, no. 2, pp. 7-12, 2022.
- C. -H. Yu, Fractional differential problem of some fractional trigonometric functions, International Journal of Interdisciplinary Research and Innovations, vol. 10, no. 4, pp. 48-53, 2022.
- C. -H. Yu, Research on two types of fractional integrals, International Journal of Electrical and Electronics Research, vol. 10, no. 4, pp. 33-37, 2022.