

**THERAPEUTIC POTENTIALS OF ALLIUM CEPA AND
CURCUMA LONGA AGAINST 2,3,7,8-
TETRACHLORODIBENZO-P-DIOXIN-INDUCED OXIDATIVE
STRESS AND LIPID IMBALANCE IN RATS**

Muhammad Sani Abdullahi

Department of Biochemistry, Faculty of Science, Ahmadu Bello University, Zaria, Nigeria.

Abstract: Industrial and environmental activities continuously release chemicals such as 2,3,7,8-tetrachlorodibenzo-p-dioxin (TCDD) that are hazardous to human health. This research seeks to assess the potentials of *Allium cepa* and *Curcuma longa* in suppressing 2,3,7,8-tetrachlorodibenzo-p-dioxin toxicity. Forty (40) male Wistar rats weighing 121.77 ± 16.23 g was divided into eight groups of five (5) rats per group. Positive control was fed standard diet, and given $2 \mu\text{g/kg/week}$ of TCDD in 0.5ml DMSO while normal control was fed standard diet and given 0.5ml dimethyl sulfoxide (DMSO)/kg body weight/week instead of TCDD. Other groups received $2 \mu\text{g/kg/week}$ of TCDD in 0.5ml DMSO and were fed varied inclusion levels of powdered *Allium cepa* and *Curcuma longa*. On the 35th day of the experimental period, the rats were sacrificed and serum prepared from collected blood for antioxidant enzymes, malondialdehyde (MDA) and lipid profile determinations. All treated groups had significantly ($p < 0.05$) higher activities of antioxidant enzymes when compared with the positive control. Malondialdehyde concentration was significantly ($p < 0.05$) lower in the treated groups than the positive control (187.87 ± 5.65 nmol/l). Lipid profiles were significantly improved upon by the inclusion of *Allium cepa* and *Curcuma longa*. Therefore, we conclude that dyslipidaemia and oxidative stress due to exposure to dioxin in rats could be suppressed by consumption of 2.5% to 5% inclusion levels of powdered onion and/or turmeric in the diet.

Keywords: 2,3,7,8-tetrachlorodibenzo-p-dioxin; glutathione peroxidase; glutathione reductase; malondialdehyde; *Allium cepa*; *Curcuma longa*.

INTRODUCTION

Dioxins are generic term for polychlorinated dibenzodioxins and polychlorinated dibenzofurans. They are among the persistent organic pollutants as their chemical stability and absorbability combine to make them last long in the body (WHO, 2016). Dioxins do bioaccumulate particularly in adipocytes. They can also bio-magnify and move long distances in the food chains (Letcher *et al.*, 2009; Shaw *et al.*, 2014). They are very toxic, the most toxic of which is 2,3,7,8-tetrachloro-dibenzo-p-dioxin (TCDD) (Ciftci *et al.*, 2018; Natalija *et al.*, 2010). They exert their toxicity by activating the Ah-receptor in the cytoplasm of vertebrates. The activation could cause compromised liver function, poor sensory abilities and distorted cognitive development (Chen *et al.*, 2006). Although known for their carcinogenicity, dioxins also mediate lipid peroxidation, cell membrane disruption,

DNA and protein damage, atherosclerosis, hypertension and other related disorders in humans via oxidative stress (Natalija *et al.*, 2010; Sally and Linda, 2009). According to WHO (2016), the lifetime tolerable number of dioxins is 70 pg/kg body weight/per month while acceptable dietary intake according to Van den Berg *et al.* (1998) is 4 pg TEQ/kg body weight/day.

The sources of dioxin are multidimensional but are mainly related to improper burning of wastes, industrial activities and leakages from power plants and obsolete stockpiles of pesticides (Patrick *et al.*, 2019). Human exposures to this deadly chemical are mostly occupational and through the environment as well as through consumption of animal products in which dioxins are stored. Clay products are high in dioxins and mothers in some African nations get exposed as they are used to consuming clay during pregnancy (Reeuwijk *et al.*, 2013). Since dioxins exert their lethal effects through oxidative stress (Hassoun *et al.*, 2006), antioxidant-enriched diets could be helpful in attenuating the effects. Turmeric has been widely studied for its function as antioxidant, which is attributed to curcumin, its major bio-active component (Priyanka *et al.*, 2017). Onions have been reported to possess antioxidant activities due to their flavonoids, anthocyanins and flavonols contents (Ana *et al.*, 2021). To the best of our knowledge, there is little information with regards to fortification of diets with turmeric and/or onion as a safe and natural means of suppressing dioxin-induced toxicity. The present study investigated the antidyslipidemic and oxidative balancing properties of diets fortified with powdered turmeric and/or onion in Wistar rats induced with dioxin toxicity.

MATERIALS AND METHODS Feed Ingredients

Soy bean meal (SBM), corn oil, palm oil, bone meal, cellulose, salt mix, methionine and pre mix were purchased from Dutsin-Ma central market. Corn starch was prepared from corn after soaking in water for 48 hours.

Identification of spices

Turmeric and onion were collected from gardens located in Ajiwa, Rimi local government area of Katsina State in Nigeria and identified as *Curcuma longa* and *Allium cepa* respectively at Department of Biological Sciences, Federal University Dutsin-Ma Katsina State. FUDMA/PSB/00143 and FUDMA/PSB/0089 voucher numbers were respectively assigned to the plants.

Chemicals

Dioxin of the type 2,3,7,8-tetrachlorodibenzo-p-dioxin was obtained from LOBA CHEMIE PVT Ltd while assay kits were purchased from reputable vendor and were products of Randox Ltd.

Preparation of Spices and Feed Formulation Onion and turmeric were dried in an envelope exposed to sunlight, pulverized and sieved using 0.05 mm sieve. The sieved samples were then mixed thoroughly in equal ratio to prepare mixed spices. Standard rodent diet was prepared by appropriately mixing corn starch, SBM, cellulose, salt, vitamin mix and mineral mix in accordance with the method of Idoko *et al.* (2022) as shown in Table 1.

Table 1: Standard rat chow

FEED COMPONENT	CONTROL DIET (g/kg)
n starch	554.5
SBM	320
Cellulose	45
Bone Meal	12.5
Palm oil	60
Salt mix	3
Vitamin/mineral mix	2.5
Methionine	2.5

Mineral mix (g/kg): CaCO₃ (15.258), CoCl₂.6H₂O (0.001), ZnCl₂ (0.001), CuSO₄.5H₂O (0.019), FeSO₄.7H₂O (1.078), MgSO₄ (2.929), MnSO₄.2H₂O (0.178), KI (0.032), KH₂PO₄ (15.559) and NaCl (5.573) Vitamin mix (g/kg diets): thiamine (0.02), riboflavin (0.03), pyridoxine (0.01), P-aminobenzoic acid (0.20), myo-inositol (2.00), biotin

(0.001), menadione (0.01), ergocalciferol (0.4), cholineHCl (2.0), and cellulose (3.31), α -tocopherol acetate (50), retinal palmitate (0.4), calcium pantothenate (0.0016) and folic acid (0.0002).

Supplemented Diet Formulation

The supplemented diets were formulated by mixing the formulated rat chow with each of the prepared spices in the ratio 95:5 and 97.5 to get 5% and 2.5% inclusion levels respectively. For 2.5% inclusion levels of the spices, the standard diet was mixed with each of the spices in the ratio of 97.5:2.5.

Preparation of Dioxin Solution

In 0.5 ml dimethyl sulfoxide, 2 μ g of 2,3,7,8-tetrachlorodibenzo-p-dioxin (TCDD) dioxin was added and allowed to dissolve completely by stirring gently. The solution was administered orally to the rats using the following relationship: $VT = BW \times 0.5$.

Where VT = volume of dioxin solution received per kg body weight; BW = body weight of individual rats.

Experimental Design

Forty (40) male Wistar rats weighing 121.77 ± 16.23 g were divided into eight (8) groups of five (5) rats per group as follow:

Group 1: Normal control was fed standard diet and given 0.5ml DMSO/week.

Group 2: Positive control was fed standard diet and given 2 μ g/kg/week of TCDD in 0.5ml DMSO. Group 3: Rats on 2.5% onion supplemented diet, and given 2 μ g/kg/week of TCDD in 0.5ml DMSO. Group 4: Rats on 5% onion supplemented diet, and given 2 μ g/kg/week of TCDD in 0.5ml DMSO. Group 5: Rats on 2.5% turmeric supplemented diet, and given 2 μ g/kg/week of TCDD in 0.5ml DMSO. Group 6: Rats on 5% turmeric supplemented diet, and given 2 μ g/kg/week of TCDD in 0.5ml DMSO (5T). Group 7: Rats on 2.5% mixture of turmeric and onion supplemented diet, and given 2 μ g/kg/week of TCDD in 0.5ml DMSO.

Group 8: Rats on 5% mixture of turmeric and onion supplemented diet, and given 2 μ g/kg/week of TCDD in 0.5ml DMSO.

All the administrations of the dioxin solution were by gavage. The rats were maintained on their respective diets *ad libitum* for a period of five (5) weeks.

Blood Sample Collection

At the end of the 5-week treatment period, the rats were weighed then anaesthetized followed by cutting of the jugular vein. Serum was obtained from collected blood samples by centrifugation at 1500 g for 15 minutes and stored at -20°C until required (Suragani *et al.*, 2014).

Determination of Indices of Dyslipidaemia

The serum concentrations of total cholesterol (TC), triglyceride (TG) and high-density lipoprotein (HDL) concentrations were estimated using commercial Randox assay kits according to manufactures instruction. Low Density Lipoprotein (LDL) was computed as described by

Friedewald *et al.* (1972): $\text{LDL} = \text{total cholesterol (Triglycerides/2.2)} - \text{HDL}$, while atherogenic index (AI) and coronary risk index (CRI) were determined following the method of Kazemi *et al.* (2018).

Atherogenic Index (AI) = LDL-C/HDL-C ; Coronary risk index (CRI) = (TC/HDL-C) .

Determination of Oxidative Status

The serum concentration of superoxide dismutase (SOD), glutathione peroxidase (GPx), glutathione reductase and catalase were determined using commercial Randox assay kits, while MDA concentrations were estimated by measuring TBARS.

Statistical Analysis

Results are presented as mean $\pm$ standard error of mean (SEM) of 3 determinations. Analysis of variance (ANOVA) and Duncan's new multiple range test were performed using SPSS version 16 (IBM Corp.) software package at 95% confidence interval.

RESULTS

The results of the lipid profile and atherogenic indices (Table 2) shows that the group exposed to TDDC but fed 5% mixed spices supplemented diet (group 8) had significantly ($p < 0.05$) least concentration of cholesterol. With the exception of group exposed to TDDC and maintained on 2.5% mixed spices supplemented diet (Group 7) which had significantly ($p < 0.05$) lower triglyceride in comparison with the normal control (Group 1), all other treated groups had significantly ($p < 0.05$) higher triglyceride when compared with the normal control but significantly ($p < 0.05$) lower if compared with the positive control (Group 2). All the treated groups had significantly ($p < 0.05$) higher HDL but lower LDL, AI and CRI as compared with the positive control (Group 2). CRI = coronary risk index

The group fed non-supplemented standard diet and given 2 $\mu\text{g/kg/week}$ of TCDD in 0.5 ml DMSO (positive control) had significantly ($p < 0.05$) lower activities of all the studied antioxidant enzymes but significantly ($p < 0.05$) higher concentration of MDA compared with the Normal control. The activities of these enzymes were significantly ($p < 0.05$) higher in the treated groups when compared with the positive control. The concentration of MDA was however lower ($p < 0.05$) in these treated groups if compared with the positive control (Table 3).

DISCUSSION

Increasing atherogenic index and coronary risk index as seen in the group of rats exposed to dioxin and fed standard diet correlates with increasing atherosclerosis and myocardial infarction (Donat-Vargas *et al.*, 2020; Kazemi *et al.*, 2018; Vennila and Pugalendi, 2012) and is consistent with the reports of Natalija *et al.* (2010) and Sally and Linda (2009) suggesting that exposure to dioxin is a high-risk factor for atherosclerosis. Similarly, Ida *et al.* (2019) reported that dyslipidemia is a major risk factor in the development of cardiovascular disease. This creates a vicious circle as damage to the heart exacerbates dyslipidaemia (Upaganlawar and Balaraman, 2012).

Diet supplementation with onion and/or turmeric moved towards normalising lipid metabolism and ameliorating dyslipidaemia, and may have prevented atherosclerosis in rats exposed to dioxin toxicity as seen in the significantly lower total cholesterol, triglyceride, LDL-C, atherogenic index, coronary risk index, and higher HDL in the groups exposed to dioxin toxicity but fed the supplemented diets relative to the group exposed to dioxin toxicity but given standard diet. Curcuminoids are the major bioactive compounds in turmeric and had been reported to possess anti-cardiometabolic risk effects (Johnston *et al.*, 2017) which according to Si *et al.* (2017) involves improvement on serum lipid panels. For instance, curcumin lowers LDL-C via decreasing LDL-C receptor (Nishiyama *et al.*, 2005). Furthermore, Imam *et al.* (2022) reported improvement in insulin resistance in hyperlipidaemic rats fed turmeric supplemented diet which could also improve lipid metabolism and prevent dyslipidaemia.

Our finding on dyslipidaemic effect of onion supplemented diet in rats exposed to dioxin could result from LDL being converted to HDL via activation of lecithin-cholesterol acyltransferase as reported by Ige and Akhigbe (2013) and/or by increased excretion of bile acids or inhibition of cholesterol absorption as demonstrated by Guan *et al.* (2010). These could be attributed to its rich fibre and flavonoids (Hamazu *et al.*, 2011). Our result is consistent with the finding of Huang *et al.* (2021) who reported that supplementation of onion has beneficial effect on dyslipidemia. It is evident from our finding that diet supplementation with onion could prevent atherosclerosis, a common feature of dioxin toxicity since according to Mattiuzzi *et al.* (2020), controlling blood lipid decreases the risk of atherosclerosis and/or cardiovascular disease.

The improvement observed in the lipid and atherogenic status of the groups of dioxin-exposed rats fed diets supplemented with onion and/or turmeric could be through scavenging excess free radicals generated by dioxin as indicated by significantly higher activities of the studied antioxidant enzymes and lower MDA levels in the exposed groups fed supplemented diets. Experiments have demonstrated that dioxin particularly TCDD cause

DNA damage, peroxidation of lipids and consequently tissue damage as a result excess generation of free radicals. Free reactive oxygen species attack lipids which negatively affect lipid fluidity leading to eventual cell death. Lipid peroxidation intermediates are toxic and could affect the DNA (Dianzani and Barrera, 2008). This is one of the major mechanisms of oxidative damages to tissues. Turmeric improves insulin sensitivity in metabolically disturbed rats (Imam *et al.*, 2022) and therefore could prevent degenerations of lipid. The major biologically active components of turmeric, curcumin lowers production of free radicals, provokes the synthesis of antioxidant enzymes (Alm-Eldeen *et al.*, 2013) and prevents formation of hydroxy radicals (Anand *et al.*, 2008). Our finding is consistent with the reported hepatoprotective effects of turmeric against TCDD toxicity which according to Ciftci *et al.* (2011) was by elevating the levels of enzyme activities involved in free radical scavenging. Diet supplemented with onion also exhibited beneficial effects against oxidative stress in rats exposed to dioxin. Onion is rich in quercetin, a flavonoid seen at the centre of its biological activities. Seref *et al.* (2014) had reported onion attenuated oxidative stress in Cd-induced oxidative stress in rats. Earlier, Obioha *et al.* (2009) had postulated antioxidative properties of onion to be by sparing the consumption of SOD, GSH-Px and catalase or by forming Cd-flavonoid complex. Although the mixed spices had beneficial effects in attenuating development of dyslipidaemia and oxidative stress, it is not clear at this point if the combination is superior to the use of the individual spices in mitigating dioxin toxicity.

CONCLUSION

Exposure to TCDD causes atherosclerosis and oxidative stress in rats which could be suppressed by consumption of 2.5% to 5% inclusion levels of powdered onion and/or turmeric. However, such supplementation should be done with caution since we are not able to establish in this study if any deleterious effect is associated with such inclusion levels and therefore necessitate further investigations.

REFERENCES

- Alm-Eldeen, A. A., Mona, M. H., Shati, A. A., & ElMekkawy, H. I. (2013). Synergistic effect of black tea and curcumin in improving the hepatotoxicity induced by aflatoxin B1 in rats. *Toxicology and Industrial Health*, 31(12), 1269-1280.
- Anand, P., Thomas, S. G., & Kunnumakkara, A. B. (2008). Biological activities of curcumin and its analogues (Congeners) made by man and mother nature. *Biochemical Pharmacology*, 76(11), 1590-1611.
- Ana, V. G., Mercedes, V., Estrella, E., Ceferino, C., Marta, F., Gerardo, F. B., & Miguel, P. (2021). Flavonol composition and antioxidant activity of onions (*Allium cepa* L.) based on the development of new analytical ultrasound-assisted extraction methods. *Antioxidants*, 10(2), 273.
- Chen, H. L., Su, H. J., Guo, Y. L., Liao, P. C., Hung, C. F., & Lee, C. C. (2006). Biochemistry examinations and health disorder evaluation of Taiwanese living near incinerators and with low serum PCDD/Fs levels. *Science of the Total Environment*, 366(2-3), 538-548.
- Ciftci, O., Sefa, A. D., Basak, N. T., & Tashdere, A. (2018). Beta-glucan prevents toxic effects of 2,3,7,8-TCDD in terms of oxidative and histopathological damage in heart tissue of rats. *Brazilian Journal of Pharmaceutical Sciences*, 54(3), 1-7.
- Ciftci, O., Ozdemir, I., Tanyildizi, S., Yildiz, S., & Oguzturk, H. (2011). Antioxidative effects of curcumin, β -myrcene and 1,8-cineole against 2,3,7,8-tetrachlorodibenzo-p-dioxin-induced oxidative stress in rats' liver. *Toxicology and Industrial Health*, 27(5), 447-453.

- Dianzani, M., & Barrera, G. (2008). Pathology and physiology of lipid peroxidation and its carbonyl products. In Álvarez, S., & Evelson, P. (Eds.), *Free Radical Pathophysiology* (pp. 19-38). Transworld Research Network.
- Donat-Vargas, C., Moreno-Franco, B., Laclaustra, M., Sandoval-Insausti, H., Jarauta, E., & Guallar-Castillon, P. (2020). Exposure to dietary polychlorinated biphenyls and dioxins, and its relationship with subclinical coronary atherosclerosis: The Aragon Workers' Health Study. *Environment International*. <https://doi.org/10.1016/j.envint.2019>
- Friedewald, W. T., Levy, R. I., & Fredrickson, D. S. (1972). Estimation of low-density lipoprotein cholesterol without the use of the preparative ultracentrifuge. *Clinical Chemistry*, 18(6), 499-502.
- Guan, L., Chung, H. Y., Su, Y., Jiao, R., Peng, C., & Chen, Z. Y. (2010). Hypocholesterolemic activity of onion is mediated by enhancing excretion of fecal sterols in hamsters. *Food & Function*, 1(1), 84–89.
- Hamauzu, Y., Nosaka, T., Ito, F., Suzuki, T., Torisu, S., Hashida, M., Fukuzawa, A., Ohguchi, M., & Yamanaka, S. (2011). Physicochemical characteristics of rapidly dried onion powder and its anti-atherogenic effect on rats fed high-fat diet. *Food Chemistry*, 129(3), 810–815.
- Hassoun, E. A., Vodhanel, J., Holden, B., & Abushaban, A. (2006). The effect of ellagic acid and vitamin E succinate on antioxidant enzymes activities and glutathione levels in different brain regions of rats after subchronic exposure to TCDD. *Journal of Toxicology and Environmental Health*, 69(5), 381-393.
- Huang, W., Tang, G., Zhang, L., Tao, J., & Wei, Z. (2021). Effect of onion on blood lipid profile: A meta-analysis of randomized controlled trials. *Food Science and Nutrition*, 9(7), 3563-3572.
- Ida, S., Kaneko, R., & Murata, K. (2019). Efficacy and safety of pemafibrate administration in patients with dyslipidemia: A systematic review and meta-analysis. *Cardiovascular Diabetology*, 18(1), 38.
- Imam, N. U., Idoko, A. S., Osibemhe, M., Lawal, N., & Zaharaddeen, A. S. (2022). Obesity and insulin resistance components of metabolic syndrome induced by high-fructose diet in wistar rats could be attenuated by spices-supplemented diets. *Journal of Applied Science and Environmental Management*, 26(5), 893-901.
- Idoko, A. S., Abdullahi, A. B., Maibulangu, M., Nura, L., Imam, N. U., Bonomi, Z. M., Muhammed, F., & Umar, S. (2022). *Allium sativum* and *Curcuma longa* powder protect against hepatotoxic and nephrotoxic effects of high-fructose diet. *FUOYE Journal of Pure and Applied Sciences*, 7(1), 59-78.
- Ige, S. F., & Akhigbe, R. E. (2013). Common onion (*Allium cepa*) extract reverses cadmium-induced organ toxicity and dyslipidaemia via redox alteration in rats. *Pathophysiology*, 20(4), 269–274.
- Johnston, T. P., Korolenko, T. A., Pirro, M., & Sahebkar, A. (2017). Preventing cardiovascular heart disease: Promising nutraceutical and non-nutraceutical treatments for cholesterol management. *Pharmacological Research*, 120, 219–225.

- Kazemi, T., Hajihosseini, M., Moossavi, M., Hemmati, M., & Ziaee, M. (2018). Cardiovascular risk factors and atherogenic indices in an Iranian population: Birjand East of Iran. *Clinical Medicine Insights: Cardiology*. <https://doi.org/10.1177/1179546818759286>
- Letcher, R. J., Gebbink, W. A., Sonne, C., Born, E. W., McKinney, M. A., & Dietz, R. (2009). Bioaccumulation and biotransformation of brominated and chlorinated contaminants and their metabolites in ringed seals (*Pusa hispida*) and polar bears (*Ursus maritimus*) from East Greenland. *Environment International*, 35(8), 1118-1124.
- Mattiuzzi, C., Sanchis-Gomar, F., & Lippi, G. (2020). Worldwide burden of LDL cholesterol: Implications in cardiovascular disease. *Nutrition, Metabolism and Cardiovascular Diseases*, 30(2), 241–244.
- Nishiyama, T., Mae, T., Kishida, H., Tsukagawa, M., Mimaki, Y., Kuroda, M., et al. (2005). Curcuminoids and sesquiterpenoids in turmeric (*Curcuma longa* L.) suppress an increase in blood glucose level in type 2 diabetic KK-ay mice. *Journal of Agricultural and Food Chemistry*, 53(4), 959-963.