



DISTRIBUTION OF ABO BLOOD GROUPS AND RH FACTORS IN THE POPULATION OF AL-MUTHANNA GOVERNORATE

Zainab Mohammad Al-Khazraji

Open Educational College - Muthanna Study Centre, Al-Muthanna Governorate, Iraq

Abstract: The ABO blood groups distribution was studied and Rhesus in (9900) individual of the Samawah, Rumaitha, Al-Warka, Al-Khader, Al-Salman judiciary in the Al-Muthanna governorate. Our research's findings indicated a high incidence of group O (3373) (38.07 %) male and female. Blood group (AB) showed a low incidence (793) (8.01%) male and female. Blood group (A) show higher incidence (3118) (31.49%) male and female. Blood group (B) showed low incidence (2616) (26.42%) male and female. Rh factors showed higher incidence in this study of (Rh^{+ve}) (9327) (94.2%) comparison with very low incidence of (Rh^{-ve}) 573 (5.79%). The distribution ratio was studied (Rh^{+ve}) ratio in (9327) individual males and females, samples showed high incidence in (O^{+ve}) ratio as followed (3127) (33.53 %) While (AB^{+ve}) showed low distribution ratio in all studied samples (766) (8.21 %). Samples showed high ratio in (O^{-ve}) (246) (43.93 %), while showed low ratio in (AB^{-ve}) (27) (4.71 %). This research was done to determine (Rh) and (ABO) distribution between the whole population of AL Muthanna Governorate.

Keywords: distribution. ABO. Rh^{-ve} . Rh^{+ve} . Rh

Introduction

The two genetic markers that are most frequently studied are the ABO blood groups and (Rh) antigens. Race and location have a big impact on the phenotype and ABO and Rh genes, despite the fact that the antigens responsible for these blood groups remain constant throughout a person's lifetime (1). In 1900, Landsteiner the first discovery of a human blood groups system ABO (2).

The term blood groups refer to the (RBC) antigen and a number of genes that determine the blood group's specificity and can be allele carefully linked on the same chromosome (3). ABO blood groups differ between ethnic groups and geographical regions (4). A person almost always has the same blood type, but very rarely does this change due to the addition of an antigen or its suppression, such as in autoimmune diseases or malignancies (5). Although their parents' genes determine their blood type, siblings' blood types vary and antigens A and blood cells (6). Blood type A is the result of having the (A) antigen, blood type B is the result of having the (B) antigen, blood type AB is the result of having both the (A,B) antigens and blood group O is the result of not having the (A,B) antigens (7). The ABO blood groups are regulated by one gene site located on chromosome nine of the autosomal genome (8). Depending on Rh antigens are present or absent on the surface of red blood cells,

blood groups in the Rh pairs of closely related allelic genes regulate the synthesis of the Rh antigen (9;10). The entire human population involved the same ABO and rhesus systems, despite the fact that they have different frequencies and distributions of particular types in various racial, ethnic and socioeconomic groups or among various populations (11).

The ABO gene produces the enzyme glycosyltransferase gene variations cause enzyme polymorphisms which affect the enzyme attaches N-acetylgalactosamine (Ag-A), Galactose (Ag-B), or no sugar (type O) (12).




Figure (1): ABO antigen privacy only the carbohydrate portion of the antigen is shown, with the (ABO) antigens differing by one sugar at the antigen end (13).

Materials & Methods

This study was conducted in 2022 over the course of six months, the study population include (9900) people a good health (who want to get married) 4950 were Male, and 4950 were Female randomly selected from resident people Samawah, Rumaitha, Al-Warka, Al-Khader, Al-Salman judiciary in the Al-Muthanna governorate. Blood groups were specified based on the agglutination using ABO and Rh blood group tests that have been performed by (Biotec Laboratories, United Kingdom).

Statistical Analysis

Collected data was treated using the (SPSS version 17). Descriptive statistics included statistical tables of numbers and percentages.

Results

The current study included (9900) samples to locate the distribution of (ABO) blood groups and (Rh), (Table 1) show high incidence of group O (3373) (34.07 %) male and female. While blood group (AB) showed low (793) (8.01%) in male and female, group A (3118) (31.49%) and group B (2616) (26.42%).

(Table 1): distribution of ABO blood groups system.

Blood groups	Male	Female	Total	Percent
A	1578	1540	3118	31.49 %
B	1312	1304	2616	26.42 %

AB	394	399	793	8.01 %
O	1666	1707	3373	38.07 %
Total	4950	4950	9900	

(Table 2) show high incidence of group **O⁺ve** (3127) (33.53 %) male and female. While blood group **AB⁺ve** showed low incidence in the city as followed (766) (8.21 %) male and female. Group **A⁺ve** (2954) (31.67%) and group **B⁺ve** (2480) (26.59%).

(Table 2): distribution of **Rh⁺ve** blood groups system.

Blood groups	Male	Female	Total	Percent
A⁺ve	1493	1461	2954	31.67 %
B⁺ve	1248	1232	2480	26.59 %
AB⁺ve	384	382	766	8.21 %
O⁺ve	1547	1580	3127	33.53 %
Total	4672	4655	9327	

(Table 3) show high incidence of group **O⁻ve** (246) (43.93 %) male and female. While blood group **AB⁻ve** showed low incidence in the city as followed (27) (4.71 %) male and female.

Group **A⁻ve** (164) (28.62%) and **B⁻ve** (136) (23.73%).

(Table 3): distribution of **Rh⁻ve** blood groups system.

Blood groups	Male	Female	Total	Percent
A⁻ve	85	79	164	28.62 %
B⁻ve	64	72	136	23.73 %
AB⁻ve	10	17	27	4.71 %
o⁻ve	119	127	246	43.93 %
Total	278	295	573	

Table (4) shows the distribution of ABO and Rh blood groups across the Al-Muthanna governorate's geographic regions. blood type among the five cities in our study. The pattern that was seen was as follows ($O > A > B > AB$). Among the five cities and **Rh⁺ve** was more prevalent than **Rh⁻ve**.

(Table 4): Distribution of ABO and Rh blood groups in zones Al-Muthanna governorate.

Blood groups	A	B	AB	O	Rh ⁺ ve	Rh ⁻ ve
Zones						
Samawah 3663	1132 (30.90%)	982 (26.81%)	142 (3.88%)	1407 (38.41%)	3419 (94.34%)	217 (5.66%)
Imaitha 2574	743 (28.87%)	616 (23.93%)	205 (7.96%)	1010 (39.24%)	2299 (89.32%)	275 (10.68%)
Warka 1980	642 (32.43%)	498 (25.15%)	89 (4.49%)	751 (37.93%)	1837 (92.78%)	143 (7.22%)
Khader 1287	372 (28.91%)	307 (23.85%)	109 (8.47%)	499 (38.77%)	1185 (92.07%)	102 (7.93%)
Salman 396	132	93	21 (5.31%)	150	370	26

	(33.33%)	(23.48%)		(37.88%)	(93.43%)	(6.57%)
--	----------	----------	--	----------	----------	---------

Discussion

The blood group AB had the lowest prevalence (8.01%) and the blood group O had the highest (38.07%) in this study. This study is consistent with earlier research conducted in southern Iraq (14;15;16), central parts of Iraq (17;18;19) and north of Iraq (20), which states group AB signed up with the lowest percentage, and phenotype O showed the most hesitation. In this study, it was observed that blood group O^{+ve} is the highest (33.53 %) and AB^{+ve} was (8.21 %) table 2, o^{-ve} is the highest (43.93 %) and AB^{-ve} was (4.71 %) table 3. This study is in agreement with previous study (21). Consequently, other studies have found a similar pattern of distribution. In Northwest Ethiopia (22), Western India (23), Tanzania (24), Mogadishu, Benadir, Somalia (25), Nigeria (26).

Conclusions

According to our research, groups O and AB were the most and minimum respectively, among blood donors in the Al-Muthanna governorate. Therefore, that information would be useful to blood banks. Additionally, we want to look into how ABO blood types relate to both fertility and spontaneous recurrent abortion.

References:

- Agrawal, A., Tiwari, A. K., Mehta, N., Bhattacharya, P., Wankhede, R., Tulsiani, S., et al. (2014). ABO and Rh (D) group distribution and gene frequency: The first multicentric study in India. *Asian Journal of Transfusion Science*, 8(2), 121-125.
- Chandra, T., & Gupta, A. (2012). Frequency of ABO and rhesus blood groups in blood donors. *Asian Journal of Transfusion Science*, 6(1), 52-53.
- Legese, B., Shiferaw, M., Tamir, W., & Tiruneh, T. (2021). Distribution of ABO and rhesus blood group phenotypes among blood donors at Bahir Dar Blood Bank, Amhara, Northwest Ethiopia: A retrospective cross-sectional study. *Journal of Blood Medicine*, 12, 849-854. <https://doi.org/10.2147/JBM.S329360>
- Mourant, A. E., Kopec, A. C., & Damaniewska-Sobczak, K. (1976). Northern and central Europe. In *Distribution of human blood groups and other polymorphisms* (2nd ed., pp. 62–69). Oxford University Press.
- Kremer, H. I., Koopmans, M., de Heer, E., Bruijn, J., & Bajema, I. (2007). Change in blood group in systemic lupus erythematosus. *Lancet*, 369(9557), 186–187.
- Cooling, L. (2015). Blood groups in infection and host susceptibility. *Clinical Microbiology Reviews*, 28(3), 801-870.
- Shafie, A. H., & Mosab, N. M. (2021). Pattern of ABO and Rhesus blood group distribution among students of Jamhuriya University of Science and Technology, Mogadishu, Benadir, Somalia. *International Journal of Medical Research & Health Sciences*, 10(11), 15-18.

- Wikipedia. (2009). ABO blood group system. The free encyclopedia. Retrieved from <http://www.geneames.org/data/hghc-data.php?match=ABO>
- Talukder, S. I., & Ruhini, K. D. (2010). Distribution of ABO and Rh blood groups in Dinajpur Medical College. *Dinajpur Medical College Journal*, 3(2), 55-58.
- Behra, D., & Joshi, D. (2013). Distribution of ABO blood group and RH (D) factor in Western Rajasthan. *Journal of Medical Research*, 3(1), 73-75.
- Sidhu, S. (2003). Distribution of the ABO blood groups and Rh (D) factor among the scheduled caste population of Punjab. *Anthropologist*, 5(2), 203-204.
- Yamamoto, F., Clausen, H., White, T., Marken, J., & Hakomori, S. (1990). Molecular genetic basis of the histo blood group ABO system. *Nature*, 345(6272), 229-233.
- Goldsby, R. A., Kindt, T. J., Osborne, B. A., & Kuby, J. (2000). *Kuby immunology* (4th ed.). W. H. Freeman.
- Taha, J. Y. (2005). Frequency of various Rh antigens in Basrah Province, Iraq. *The Medical Journal of Basrah University*, 53-54.
- Al-Fartosi, K. G. (2008). The association between ABO blood group and spontaneous abortion. *Basrah Journal of Science (B)*, 26(1), 1-10.
- Al-Ali, H. S. (2008). Association of ABO and Rh blood groups with diabetes mellitus and hypertension in Basrah City. *Basrah Journal of Science (B)*, 26(1), 29-37.
- Hasan, T. A. A., & Nasir, I. A. (2016). Blood groups and their association with chronic diseases in the local population of Karbala city. *Journal of the University of Kerbala*, 14(3), 1-10.
- K. H., Salah, & Al-Maliki, A. H. (2015). Normal distribution of ABO blood group and Rhesus factor in Al-Najaf province. *European Journal of Experimental Biology*, 5(7), 18-21.
- Al-Khatieeb, M. M., Al-Joubori, S. K., & Taha, S. S. (2018). Association of ABO blood group and Rhesus factor with dental malocclusion in a population of Baghdad, Iraq. *International Journal of Medical Research & Health Sciences*, 7(1), 165-169.
- Jaff, M. S. (2010). ABO and rhesus blood group distribution in Kurds. *Journal of Blood Medicine*, 1, 143-146.
- Adeyemo, O. A., & Soboyejo, O. B. (2006). ABO and Rh blood group distribution in the African population. *African Journal of Biotechnology*, 5(22), 2062-2065.

- Biruk, L., Mikru, S., Workineh, T., & Tegenaw, T. (2021). Distribution of ABO and Rhesus blood group phenotypes among blood donors at Bahir Dar Blood Bank, Amhara, Northwest Ethiopia: A retrospective cross-sectional study. *Journal of Blood Medicine*, 12, 849-854.
- Chandekar, A. S., Amonkar, P., Desai, H., Valvi, N., & Puranik, G. (2017). Distribution of ABO and Rh-D blood groups among blood donors: Western India data. *International Journal of Scientific Study*, 5(3), 1-6.
- Jahanpour, O., Pyuza, J. J., Ntiyakunze, E. O., Mremi, A., & Shao, E. R. (2017). ABO and Rhesus blood group distribution and frequency among blood donors at Kilimanjaro Christian Medical Center, Moshi, Tanzania. *BMC Research Notes*, 10(1), 738. <https://doi.org/10.1186/s13104-017-3037-3>
- Ermış, T., Adan, N. A., Gacal, N., Noh, R. A., & Arslan, E. (2022). ABO and Rhesus blood group distribution in Mogadishu, Somalia. *Journal of Blood Medicine*, 13, 213-218.
- Egesie, U. G., Egesie, O. J., Usar, I., & Johnbull, T. O. (2008). Distribution of ABO, Rhesus blood, and haemoglobin electrophoresis among the undergraduate students of Niger Delta State University, Nigeria. *Nigerian Journal of Physiological Sciences*, 23(1-2), 5-8.