

**ANTIBACTERIAL POTENTIAL OF GINGER AND
GARLIC EXTRACTS AGAINST PATHOGENIC
BACTERIA****Fatimah Olufunke Bello**

Department of Microbiology, Federal University Gusau, Zamfara State, Nigeria

Abstract: Cases of fake/weak drugs are abundant everywhere in the country today, especially in the use of antibiotics, which leads to an increase level of drug resistance among microorganisms which were known to be susceptible to particular antibiotics. Thus, the need to test for the antibacterial activity of ginger and garlic against some pathogenic microorganisms should be employed to support its use as an ingredient for herbal medicines. This study explores the potential of ginger ethanol extracts, ginger chloroform extract, garlic ethanol extract and garlic chloroform extract tested against some clinically relevant bacterial strains. These include; *Citrobacter* sp, *Escherichia coli*, *Staphylococcus aureus* and *Pseudomonas auregenosa*, using agar well diffusion technique. Ciprofloxacin (1 mg/ml) was used as standard control. The test concentration of each extract was set as 100 mg/ml, 50 mg/ml, 25 mg/ml and 12.5 mg/ml. The phytochemical screening revealed the presence of tanins, flavonoids, saponins, cardiac glycosides, steroid, alkaloid, volatile oils, balsams and terpenoids. The results showed a higher zone of inhibition of 28.45 mm at 100 mg/ml concentration of ginger ethanol extract against *S. aureus* and garlic chloroform extract with the least activity against *Citrobacter* spp. with 6.30 mm at 100 mg/ml. The MIC of ginger ethanol extract against the test organisms was determined, which show values between 12.5 and 95 mg/ml. In comparison, the ginger chloroform extract had MIC values between 12.5 and 50 mg/ml. The garlic ethanol extract had MIC value of 50 mg/ml while the garlic chloroform extract had MIC value of 100 mg/ml. The MBC of ginger ethanol and chloroform extracts was between 12.5 and 100 mg/ml. The garlic ethanol and chloroform extracts had MBC values only at 100 mg/ml. The result of this study show that the extracts had activity against the test organisms and as such could be a potential therapeutic against the tested organisms.

Keywords: *Zingiber officinale* (Ginger); *Allium sativum* (Garlic) Phytochemicals; Antibacterial

Introduction

Antibiotic resistance has become a serious problem and affects almost every bacterial specie. Resistance to multiple antibiotics has developed among many common pathogens, such as *Staphylococcus*, *Pneumococci*, *Pseudomonas* organisms and this problem is steadily increasing worldwide (Olofsson and Cars, 2007). Around 90-95% of *Staphylococcus aureus* strains is penicillin resistance worldwide. In Asian countries, 70-80% of the

same strains are methicillin resistance (Hemaiswarya et al., 2008). Sometimes antibiotics are associated with adverse effects on host, which include depletion of beneficial gut, mucosal microorganisms, immune suppression, hypersensitivity and allergic reaction. Some drug-resistant bacteria have complicated the treatment of infectious diseases in immune compromised AIDS and cancer patients (McGaw et al., 2000). One way to beat this downside of drug resistance is by getting new molecules from natural resources.

Plants are known to produce a variety of compounds and medicinal properties to prevent infections from a wide range of microorganisms including plant pathogens and environmental organisms (Bentley, 1997; Savithramma et al., 2011; Chung et al., 2011). Therefore, alternative antimicrobials are used from botanical sources which provide flexibility and diversity. In many developing countries large portion of population depends on the traditional system of medicine to treat variety of disease (McGaw et al., 2000). The World Health Organization (WHO) reported that 80 % of world population relies chiefly on traditional medicine, which involve the medicinal plant extracts or their active constituents (Ahmad et al., 1998). Subsequently advancement of common antimicrobial agents from plant source would serve as a promising approach.

Ginger, *Zingiber officinale* (commonly called 'jinja' in Igbo, 'citta' in Hausa and 'Atale' in Yoruba). Ginger is a member of the family Zingiberaceae a small family with more than 45 genera and 800 species, its scientific name is *Zingiber officinale* (Foster, 2011). Ginger is an erect, slim herbaceous perennial plant growing from one to three feet in height. Its stem is surrounded by the sheath bases of the two ranked leaves. A club like spike of yellowish, purple lipped flowers, it has a greenish yellow brick with rarely flowers in cultivation (Tyler, 2002). It possesses a fleshy and thick underground rhizome and having one or more aerial leafy stems that grows up to 1.25m tall. Ginger is grown in the tropical weathers of Australia, West Africa, India, Jamaica, Brazil, China, and some parts of the United States (Suruchi et al., 2016). In the first year of growth, it produces a green straight stalk like stem about 60 cm high growing from the rhizome. Its leaves grow and measures about 12-30 cm long which dies off each year. The crop grows preferably in warm, sunny conditions, and may profit from shade during hot days, especially when young. Shading is however generally considered redundant. The optimum rainfall is 2500-3000 mm, well distributed over the year. (Shubha, 2015). Ginger plays an important medicinal role due to the presence of certain constituents such as gingerol, paradol, shogaol, zingerone, zerumbone, terpenoids and ginger flavonoids (Arshad et al., 2014).

Garlic, *Allium sativum* (commonly called 'aayu' in Yoruba, 'ayo-ishi' in Igbo and 'tafarnua' in Hausa), is a perennial bulbous plant that initially came from middle Asia and is at present grown globally. It belongs to the family Alliaceae. Garlic can grow up to 2 feet in height or more. The bulb is the main part of the plant which is used for medicine (Steven, 2015). Each garlic bulb is made up of 4 to 20 cloves. Each garlic clove may weigh about 1 gram in weight. Fresh, aged, or dried garlic can be used as garlic supplements. Each of the supplements may have different effects to the body (Sethi et al., 2014). It is commonly used as seasoning. Its close relatives include the onion, shallot and leek. The head of garlic (the most commonly used plant part) comprises numerous discrete cloves whereas the leaves and stems are sometimes eaten, particularly while immature and tender. The medicinal potency of garlic is due to glycoside, vitamin B, C, and D allisatin II and I. It also contains volatile sulphur oil, which has a vermifugal action (Arshad et al., 2014). It has been used throughout recorded history for both culinary and medicinal purposes. It has a characteristic pungent, hot, flavour that mellows and sweetens considerably with cooking. Its typical pungent odour antibacterial activity depends on allicin, which is produced by enzymatic (alliin-lyase) hydrolysis of alliin after cutting and crushing of the cloves (Onyeagba et al., 2004)

Methods Sample Collection and Processing

The plants materials (Garlic and Ginger) were obtained from Kabuga market, Kano, Kano State, Nigeria. They were placed in separate polythene bags and transported immediately to the laboratory of the Department of Microbiology, Bayero University, Kano. The samples were identified and authenticated by comparing them with known samples. the plant materials were washed with clean water to get rid of sand particles. They were chopped and partially allowed to air-dry in the shade at room temperature for five days in order to remove excess moisture. The garlic bulbs were separated into cloves. The cloves skins were peeled off and the cloves were sliced and also air dried at ambient temperature for about four weeks. After drying, pieces of *Allium sativum* and *Zingiber officinale* were grinded to fine particles each, utilizing a suitable sterile electric blender to obtain a homogenous sample.

Preparations of Extract

25 g of the powdered samples each was extracted with 250 mls of ethanol and chloroform by cold maceration method as described by Handa et al. (2008); Jolly et al. (2022), with some slight modifications. The containers were left at 25 °C for 4 days (96 hours). The suspensions were filtered using Whatman no.1 filter paper. The filtrates were concentrated at 90 °C using water bath and delivered into sterile clean containers with suitable labeling and were kept at 4°C in a refrigerator until further use. The percentage yield of each extract obtained was calculated using the formula.

$$We/Wp \times 100$$

Where; *We*= weight of the extract,

Wp = weight of the powdered material used for the extraction.

Sterility of the Extracts

One (1) ml of the extracts was added separately in two test tubes containing 5 ml of sterile nutrient broth. They were incubated at 37 °C for 24 hours. The extracts were cleared after incubation indicating the absence of contaminant which could have caused a turbid appearance in the tube (Iotsor et al., 2019).

Qualitative Phytochemical Screening

Phytochemical screening of the extracts were conducted qualitatively as stated by Harborne (1973); Yusha'u et al. (2009); Tafinta et al. (2020). Tanins, flavonoids, saponins, cardiac glycosides, reducing sugars, steroid, alkaloid, volatile oils, balsams and terpenoids were carried out accordingly.

Test Microorganisms

The test organisms were collected from Microbiology Department Bayero University Kano. The isolates were subjected to Gram's staining and other biochemical tests according to standard procedures and identified as *Staphylococcus aureus* (MRSA), *Escherichia coli*, *Pseudomonas auregenos* and *Citrobacter* sp. (Clarke and Cowan, 1952; Daniel, 2000). The pure isolates were then stored on Nutrient agar slant bottles at a temperature of 4°C until further use.

Preparation of extract concentration

The concentration was prepared by dissolving 0.1 g of crude extract into one milliliter of Dimethylsulphuroxide (DMSO) in a clean grease free vial bottle, to obtain a stock of 100mg/ml concentration. Using double serial dilution methods the following concentration of 50mg/ml, 25mg/ml and 12.5 mg/ml were made. 1mg/ml of ciprofloxacin was used as positive control (Afolabi *et al.*, 2020).

Antibacterial Assay

The agar well diffusion method was used to investigate the antimicrobial properties of the extracts as described by NCCLS (2000); Schumacher *et al.* (2018). All media were prepared and sterilized according to manufacturer's

instructions. Within 15 minutes of adjusting the turbidity of the inocula suspension, a sterile swab stick was used to inoculate the inocula onto dried surface of sterile prepared Mueller Hinton agar plates. In each case, streaking was repeated two more times, rotating the plate approximately 60° each time to ensure an even distribution of the inoculums. The inoculated plates were allowed to stay for about 3-5 minutes for the surface of the agar to air dry. While the plates were drying, four various concentrations of the extracts were prepared. A sterilized cork borer of an internal diameter of 4mm was then used to punch five holes in the inoculated medium, the bottom of the wells were then closed using 1 ml of sterile Mueller Hinton agar and the various concentrations of the Prepared extracts were dispensed into the respective labeled holes (100 mg/ml, 50 mg/ml, 25 mg/ml and 12.5 mg/ml). 1 mg/ml of Ciprofloxacin was used as positive control. Four plates were made for each test organism and the procedure was repeated for the other organisms. The plates were kept in refrigerator for about 4 - 5 hours for complete diffusion of the extracts and incubated at 37°C for 24 - 48 hours. After the incubation period, the diameter of each zone of inhibition was measured with a handheld vernier caliper in mm and the results were recorded.

Determination of Minimum inhibitory concentration (MIC)

The results of antibacterial activity were used for the determination of MIC. Using single serial dilution method using dimethylsulphuroxide (DMSO) to obtain a stock of 200mg/ml. the following concentrations of 100 mg/ml, 95 mg/ml, 90 mg/ml and 85 mg/ml were prepared but depending on the least concentration that had reasonable activity against the test organisms. If reasonable activity starts from 50mg/ml then the MIC concentrations starts from 50, 45, 40 and 35 mg/ml. The test organism was standardized using 0.5 McFarland standards. Six test tubes were used, each test tube was leveled according to concentration while the last two test tubes serves as positive and negative control. 2ml of nutrient broth was dispensed into sterile test tubes and autoclaved at 121°C for 15 min. the following volume (1 ml, 0.95 ml, 0.9 ml, and 0.85 ml) were removed from the autoclaved nutrient broth and replace with appropriate concentration from the stock solution. One hundred micro liter (0.1 ml) of standardized organisms were added into five test tubes respectively. Tubes containing broth and extracts serves as positive control while tubes containing broth and inocula serves as negative control. This procedure was repeated for the remaining extracts and test organisms. The tubes were observed after 24 hours of incubation to determine minimum inhibitory concentration, i.e the lowest concentration that showed no evidence of growth or turbidity (Yusha'u *et al.*, 2009).

Determination of Minimum Bactericidal Concentration (MBC)

Mueller-Hinton agar was prepared and the petri plates were separately inoculated with sample from each of the test tubes that showed no evidence of growth. The plates were further incubated at 35 °C for 24 hours and observed. The least concentration at which the organism did not grow was taken as the minimum bactericidal concentration. (Iotsor *et al.*, 2019; Yusha'u *et al.*, 2009).

RESULTS

Table 1.0 shows the physical properties of Ginger and Garlic Extracts, ginger extract was recovered as a dark brown, loose substance that smelled spicy-sweet while the garlic extract was a golden-yellow, sticky residue with a harsh disagreeable scent. The percentage yield of extract show that ginger ethanol extract yielded the highest amount of 10 %, followed by ginger chloroform extract with 8.4 %, followed by garlic ethanol extract with 3.6 % and garlic chloroform extract with 2.4 %.

Table 1: Physical Properties of the Plant Extracts

Characteristics	ETH. ginger	CCl4 ginger	ETH. garlic	CCl4 garlic
Weight	25g	25g	25g	25g
Odour	Spicy sweet	Spicy sweet	Harsh scent	Hash scent
Volume of solvent	250ml	250ml	250ml	250ml
Weight of extract yield	2.5g	2.1g	0.9g	0.6g

Percentage yield 10% 8.4% 3.6% 2.4%

Key: ETH.ginger = Gnger ethanol extract, ETH.garlic= Garlic ethanol extract, CCl4.ginger = Ginger chloroform extract, CCl4.garlic = Garlic chloroform extract

Table 2: Phytochemical Constituents Found in Ginger and Garlic Extracts

Phytochemical	ETH. Ginger	CCl4 ginger	ETH. garlic	CCl4 garlic
Tanins	-	-	+	-
Flavonoids	+	+	-	-
Saponins	+	+	+	-
Cardiac glycosides	+	+	+	+
Reducing sugas	-	-	-	-
Steroids	+	+	-	-
Alkaloids	+	-	+	-
Volatile oil	+	+	+	+
Balsams	+	+	-	-
Terpenoids	+	+	-	-

Key: ETH.ginger = Gnger ethanol extract, ETH.garlic= Garlic ethanol extract, CCl4.ginger = Ginger chloroform extract, CCl4.garlic = Garlic chloroform extract, + = present and - = absent.

Table 3: Zone of inhibition for the ginger ethanol extract, measured in millimeters

Test Organism	Concentration/zone inhibition				
	100mg/ml	50mg/ml	25mg/ml	12.5mg/ml	Control
Citrobacter spp.	17.15	7.97	4.00	4.00	33.18
E. coli	4.00	4.00	4.00	4.00	4.00
S. aureus	28.45	28.00	27.12	26.51	4.00
P. aeruginosa	14.03	12.84	7.27	6.99	10.21

Key: Diameter of cork borer is 4.00 mm

Table 4: Zone of inhibition for the ginger chloroform extract, measured in millimeters

Concentration/zone inhibition					
-------------------------------	--	--	--	--	--

Test Organism	100mg/ml	50mg/ml	25mg/ml	12.5mg/ml	Control
Citrobacter spp.	20.87	15.28	10.77	6.62	30.60
E. coli	4.00	4.00	4.00	4.00	4.00
S. aureus	25.98	22.78	21.58	15.79	4.00
P. aeruginosa	7.55	7.11	4.00	4.00	6.65

Key: Diameter of cork borer is 4.00 mm

Table 5: Zone of inhibition for the garlic ethanol extract, measured in millimeters

Concentration/zone inhibition					
Test Organism	100mg/ml	50mg/ml	25mg/ml	12.5mg/ml	Control
Citrobacter	4.00	4.00	4.00	4.00	30.60 spp.
E. coli	4.00	4.00	4.00	4.00	4.00
S. aureus	16.60	13.59	4.00	4.00	4.00
P. aeruginosa	4.00	4.00	4.00	4.00	4.19

Key: Diameter of cork borer is 4.00 mm

Table 6: Zone of inhibition for garlic chloroform extract, measured in millimeters

Concentration/zone inhibition					
Test Organism	100mg/ml	50mg/ml	25mg/ml	12.5mg/ml	Control
Citrobacter spp.	6.30	4.00	4.00	4.00	25.19
E. coli	4.00	4.00	4.00	4.00	4.00
S. aureus	4.00	4.00	4.00	4.00	4.00
P. aeruginosa	5.63	4.00	4.00	4.00	15.21

Key: Diameter of cork borer is 4.00 mm

Table 7: Minimum Inhibitory Concentration (MIC) of ethanol and chloroform Extracts of ginger and garlic on test organisms.

Concentration (mg/ml)				
Test Organism	ETH. Ginger	CCl4 ginger	ETH. garlic	CCl4 garlic
Citrobacter spp.	95	25	ND	100

E. coli	ND	ND	ND	ND
S. aureus	12.5	12.5	50	ND
P. aeruginosa	45	50	ND	100

Key: ETH.ginger = Ginger ethanol extract, ETH.garlic= Garlic ethanol extract, CCl₄.ginger = Ginger chloroform extract, CCl₄.garlic = Garlic chloroform extract and ND = Not detected.

Table 8: Minimum bactericidal Concentration (MBC) of ethanol and chloroform extract of ginger and garlic extracts.

Test Organism	Concentration (mg/ml)			
	ETH. Ginger	CCl ₄ ginger	ETH. garlic	CCl ₄ garlic
Citrobacter spp.	95	50	ND	100
E. coli	ND	ND	ND	ND
S. aureus	12.5	12.5	100	ND
P. aeruginosa	90	100	ND	100

Key: ETH.ginger = Ginger ethanol extract, ETH. garlic= Garlic ethanol extract, CCl₄.ginger = Ginger chloroform extract, CCl₄.garlic = Garlic chloroform extract and ND = Not detected.

DISCUSSION

The findings of this study show that solvents with high polarity yielded higher bioactive compounds. This work correlate with the findings of Wolde *et al.* (2018) who reveals that the higher the polarity of a solvent, the more amount of bioactive compound to be obtained.

The outcomes of the phytochemical screening of ethanol and chloroform extracts of ginger and garlic indicated that majority of the phytochemicals were slightly present only in the ethanol and chloroform extract of garlic, but they were abundantly found in the ethanol and chloroform extract of ginger. Only garlic ethanol extracts contain tannins. The study findings show that the ginger ethanol and chloroform extracts contain flavonoid, which is known to act as a potent barrier against bacterial infection (Namadina *et al.*, 2021). Both the ginger ethanol and chloroform extracts contain saponin, while only the garlic ethanol extract did. High levels of saponin was shown to have structure-dependent therapeutic effects (Ayoola *et al.*, 2008). The ginger and garlic ethanol and chloroform extracts contain cardiac glycosides and volatile oil content. There was no reducing sugar in any of the extracts. Both the ginger ethanol and chloroform extracts contain steroid. It is well known that plant-derived steroids have cardiogenic effects in addition to having antibacterial and insecticidal capabilities (Alexei *et al.*, 2001). The ethanol extract of ginger and garlic contain alkaloid. Many alkaloids derived from medicinal plants exhibit biological activities such as antimicrobial, cytotoxic and pharmacological effects (Dua *et al.*, 2013; Benbott *et al.*, 2012). Terpenoids were detected in both the ethanol and chloroform extracts of ginger, whereas balsams were also discovered in both.

In the present investigation, the ginger extracts exhibited high degree of inhibitory activity against most of the 4 isolates with the exception of *E. coli* followed by the garlic extracts which showed less activity at only higher

concentrations. Among the test organisms, *Staphylococcus aureus* and *Citrobacter* sp. were most susceptible then followed by *P. aeruginosa* which show the least susceptibility to garlic chloroform extract. *E. coli* was found to be resistant to all the four extracts and this could be related to the fact that the lipopolysaccharide (LPS) layer of gram-negative bacteria in outer membrane have a high hydrophobicity which acts proly as a strong barrier against the bioactive molecules. Certain molecules can pass through cell wall of Gram-positive bacteria easier than the gram-negative bacteria because cell wall of the gram-positive bacteria contained only peptidoglycan (Ababutain, 2011). Similar finding was obtained from other researchers (Akrayi, 2014; Ababutain, 2011; Keskin & Toroglu, 2011; Nanasombat & Lohasupthawee, 2011), where they found that so many extracts of spices and herbs did not have antibacterial activity against *E. coli* tested in their studies.

The least activity of garlic observed in this study was in disagreement with earlier reports that garlic is highly effective against microorganisms (Belguith *et al.*, 2010; Yin *et al.*, 2002; Bakht *et al.*, 2011; Iwalokun *et al.*, 2004; O’Gara *et al.*, 2000). The resistivity might be as a result of the heat applied during the evaporating phase of the extracts filtrates in a water bath which may have caused the denaturing of some bioactive compounds in the garlic used. According to Gupta and Ravishankar (2005), commercial garlic showed antimicrobial activity only at 4 °C and 8 °C, indicating that antimicrobial activity of garlic is temperature dependent. Apart from temperature it is also believed that geographical location of a plant, temperature, and seasonal variation of an area may have influence over the yield of medicinal plants (Akrayi, 2014). Hence, the low or no inhibition zones observed.

However, the study findings demonstrated that compared with all the extracts and ciprofloxacin as the control drug had higher zone of inhibition producing to 30 mm against all the test isolates with the exception of *Staphylococcus aureus* that showed resistance to Ciprofloxacin. Ginger ethanol and chloroform extracts performed better than the Ciprofloxacin on *Staphylococcus aureus*.

Finally, the susceptibility screening of *Citrobacter* spp, *Escherichia coli*, *Staphylococcus aureus* and *pseudomonas auregenosa*, on the extracts were further evaluated in order to determine the minimum inhibitory concentration (MIC) and minimum bactericidal concentration (MBC) all the result obtained are within the range of 12.5mg/ml to 100mg/ml.

Conclusion

This study found that the ethanol and chloroform extract of ginger (*Zingiber officinale*) displayed strong inhibitory action against *Staphylococcus aureus* than ethanol and chloroform extract of garlic. The ginger could be regarded as a potential antibacterial agent with therapeutic potential in the treatment of bacterial infection especially those caused by *Staphylococcus aureus*.

Authors’ Contributions

Yero I.H. conceived the idea and wrote the original draft of manuscript and was also involved particularly in Protocol development, Collection of Ginger and Garlic sample and scientific identification, Aminu A.I. involved in review of the manuscript and experimental design, Ishaq

S.A. assisted in result collation and data analysis. All author’s reviewed and edited the final version of the manuscript.

Acknowledgements

The authors would like to appreciate the Bayero University Kano Microbiology laboratory technologists for their undiluted support during and after the research was completed.

Conflict of Interest

The authors declare no conflict of interest exist

Reference

- Ababutain, I. M. (2001). Antimicrobial activity of ethanolic extracts from some medicinal plants. *Australian Journal of Basic and Applied Sciences*, 5(2), 678-683.
- Afolabi, B. T., Agu, G. C., & Onajobi, I. B. (2020). Phytochemical screening and antibacterial activity of *Garcinia kola* (Hackel) and *Cola nitida* (Vent) extracts. *Nigerian Journal of Technology (NIJOTECH)*, 39(2), 379-385.
- Ahmad, I., Mehmood, Z., & Mohammad, F. (1998). Screening of some Indian medicinal plants for their antimicrobial properties. *Journal of Ethnopharmacology*, 62, 183-193.
- Akrayi, F. S. (2014). Antibacterial effect of aqueous extracts of spices and herbs against bacteria isolated from frozen meat. *Medical Journal of Islamic World Academy of Sciences*, 22(1), 30-35.
- Alexei, Y. B., Joseph, I. S., & Olga, V. F. (2009). Endogenous cardiotonic steroids: Physiology, pharmacology and novel therapeutic targets. *Pharmacological Reviews*, 61, 9–38.
- Arshad, H. R., Fahad, M., Al., S., & Salah, M. A. (2014). Active ingredients of ginger as potential candidates in the prevention and treatment of diseases via modulation of biological activities. *International Journal of Physiology, Pathophysiology and Pharmacology*, 6(2), 125-136.
- Ayoola, G. A., Coker, H. A. B., Adesegun, S. A., Adepoju-Bello, A. A., Obaweya, K., Ezennia, E. C., & Atangbayila, T. O. (2008). Phytochemical screening and antioxidant activities of some selected medicinal plants used for malaria therapy in Southwestern Nigeria. *Tropical Journal of Pharmaceutical Research*, 7(3), 1019-1024.
- Bakht, J., Tayyab, M., Ali, H., Islam, A., & Shafi, M. (2011). Effect of different solvent extracted sample of *Allium sativum* (Linn) on bacteria and fungi. *African Journal of Biotechnology*, 10, 5910–5915.
- Belguith, H., Kthiri, F., Chati, A., Sofah, A. A., Hamida, J. B., & Landoulsi, A. (2010). Study of the effect of aqueous garlic extract (*Allium sativum*) on some *Salmonella* serovars isolates. *Emirates Journal of Food and Agriculture*, 22, 189–206.
- Benbott, A., Yahya, A., & Belaïdi, A. (2012). Assessment of the antibacterial activity of crude alkaloids extracted from seeds and roots of the plant *Peganum harmala*. *Journal of Natural Product Plant Resources*, 2, 568-573.
- Bentley, R. (1997). Microbial secondary metabolites play important roles in medicine and prospects discovery of new drugs. *Perspectives in Biology and Medicine*, 40, 364-394.
- Chung, P. Y., Navaratnam, P., & Chung, L. Y. (2011). Synergistic antimicrobial activity between pentacyclic triterpenoids and antibiotics against *Staphylococcus aureus* strains. *Annals of Clinical Microbiology and Antimicrobials*, 10, 1-6.

- Clarke, P. H., & Cowan, S. T. (1952). Biochemical methods for bacteriology. *Journal of General Microbiology*, 6(1-2), 187-197.
- Daniel, M. M. (2000). *Bacteriology – Haemophilus species: Baron medical microbiology*. 8th Ed. Wright's books.
- Dua, V. K., Gaurav, V., Bikram, S., Aswathy, R., Upma, B., Dau, D. A., Gupta, N. C., Sandeep, K., & Ayushi, R. (2013). Anti-malarial property of steroidal alkaloid conessine isolated from the bark of *Holarrhena antidysenterica*. *Malaria Journal*, 12, 1-6.
- Foster, S. (2011). *Ginger Zingiber officinale: Your food is your medicine*. Retrieved from <http://www.stevenfoster.com/education/monograph/ginger.html>.
- Gupta, S., & Ravishankar, S. (2005). Foodborne pathogens and disease. *Winter*, 2(4), 330-340.
- Handa, S. S., Khanuja, S. P. S., Longo, G., & Rakesh, D. D. (2008). *Extraction technologies for medicinal and aromatic plants*. 1st Ed. Trieste, Italy: Earth, Environmental and Marine Sciences and Technologies.
- Harborne, J. B. (1973). *Phytochemical methods: A guide to modern techniques of plant analysis*. 1st Ed. New York: Chapman and Hall.
- Hemaiswarya, S., Kruthiventi, A. K., & Doble, M. (2008). Synergism between natural products and antibiotics against infectious diseases. *Phytomedicine*, 15, 639-652.
- Iotsor, B. I., Iseghohi, F., Oladoja, O. E., Raji, O. R., Yusuf, Z., Oyewole, O. A., & others. (2019). Antimicrobial activities of garlic and ginger extracts on some clinical isolates. *The International Journal of Biotechnology*, 8(1), 59-65.
- Iwalokun, B. A., Ogunledun, A., Ogbolu, D. O., Bamiro, S. B., & Jimi-Omojola, J. (2004). In vitro antimicrobial properties of aqueous garlic extract against multidrug-resistant bacteria and *Candida* species from Nigeria. *Journal of Medicinal Food*, 7, 327–333.
- Jolly, O. A., Beatrice, K., Dorothy, N., & John, K. (2022). Effect of aqueous and organic solvent extraction on in-vitro antimicrobial activity of two varieties of fresh ginger (*Zingiber officinale*) and garlic (*Allium sativum*). *Heliyon*, 8, 2405-8440.
- Keskin, D., & Toroglu, S. (2011). Studies on antimicrobial activities of solvent extracts of different spices. *Journal of Environmental Biology*, 32, 251-256.
- McGaw, L. J., Jager, A. K., & Staden, J. V. (2000). Antibacterial, anthelmintic, and anti-amoebic activity in South African medicinal plants. *Journal of Ethnopharmacology*, 72, 247-263.

- Namadina, M. M., Mukhtar, A. U., Kraye, S. I., Musa, F. M., Bah, I. H., & Maitama, F. Y. (2021). Phytochemical constituents and antibacterial activity of indigenous chewing stick (*Anogeissus leiocarpus*) stem. *Bayero Journal of Pure and Applied Sciences*, 14(1), 85-94.
- National Plant Data Center, NRCS, & USDA. (2000). The plants database for *Zingiber officinale*. Baton Rouge, USA, 5(1), 70874-4490. Retrieved from <http://plants.usda.gov>.
- Nanasombat, S., & Lohasupthawee, P. (2005). Antibacterial activity of crude ethanolic extracts and essential oils of spices against *Salmonella* and other enterobacteria. *KMITL Science and Technology Journal*, 5, 527-538.
- O'Gara, E. A., Hill, D. J., & Maslin, D. J. (2000). Activities of garlic oil, garlic powder, and their diallyl constituents against *Helicobacter pylori*. *Applied and Environmental Microbiology*, 66, 2269–2273.
- Olofsson, S. K., & Cars, O. (2007). Optimizing drug exposure to minimize selection of antibiotic resistance. *Clinical Infectious Diseases*, 45, 129-136.
- Onyeagba, R., Ugbogu, O. C., Okeke, C. U., & Iroakasi, O. (2004). Studies on the antimicrobial effects of garlic (*Allium sativum* L.), ginger (*Zingiber officinale* Roscoe), and lime (*Citrus aurantifolia* L.). *African Journal of Biotechnology*, 3(4), 552-554.
- Savithramma, N., Linga, R. M., & Suhrulatha, D. (2011). Screening of medicinal plants for secondary metabolites. *Middle-East Journal of Scientific Research*, 8, 579–584.
- Schumacher, A., Vranken, T., Malhotra, A., Arts, J. J. C., & Habibovic, P. (2018). In vitro antimicrobial susceptibility testing methods: Agar dilution to 3D tissue-engineered models. *European Journal of Clinical Microbiology and Infectious Diseases*, 37(2), 187–208.
- Sethi, N., Kaura, K., Dilbaghi, N., Parle, M., & Pal, M. (2014). Garlic: A pungent wonder from nature. *International Research Journal of Pharmacy*, 5(7), 523-529.
- Shubha, S. R. (2015). Medicinal uses of ginger (*Zingiber officinale* Roscoe) improves growth and enhances immunity in aquaculture. *International Journal of Chemical Studies*, 3(2), 8387.
- Steven, D. E. (2015). Garlic. Available from <https://www.umm.edu/health/medical/atlmed/herb/garlic>.
- Suruchi, Y., Pramod, K. S., & Md, A. A. (2016). Ginger medicinal uses and benefits. *European Journal of Pharmaceutical and Medical Research*, 3(7), 127-135.
- Tafinta, I. Y., Okoye, N. H., Batagarawa, U. S., Hamma, I. I., & Abubakar, M. M. (2020). Phytochemical screening and antifungal activities of cashew (*Anacardium occidentale* Linn.) leaves extract on some fungal isolates. *Asian Plant Research Journal*, 5(3), 30-37.

- Tyler, V. E. (2002). *The honest herbal: A sensible guide to the use of herbs and related remedies*. New York: Pharmaceutical Products Press.
- Wolde, T., Kuma, H., Trueha, K., & Yabeker, A. (2018). Anti-bacterial activity of garlic extract against human pathogenic bacteria. *Journal of Pharmacovigilance*, 6, 253.
- World Health Organization. (2001). *Legal status of traditional medicine and complementary/alternative medicine: A worldwide review*. World Health Organization, Geneva, Switzerland.
- Yin, M. C., Chang, H. C., & Tsao, S. M. (2002). Inhibitory effects of aqueous garlic extract, garlic oil, and four diallyl sulphides against four enteric pathogens. *Journal of Food and Drug Analysis*, 10, 120–126.
- Yusha'u, M., Onuorah, F. C., & Murtala, Y. (2009). In-vitro sensitivity pattern of some urinary tract isolates to *Carica papaya* extracts. *Bayero Journal of Pure and Applied Sciences*, 2(2), 75-78.